

**Warm-up:**

1. Recall the definition of an infinite improper integral:

$$\int_a^\infty f(x)dx = \lim_{t \rightarrow \infty} \int_a^t f(x)dx$$

Determine whether or not the following integral converges, and if it does, evaluate the integral.

$$\int_e^\infty \frac{1}{x(\ln x)^2} dx$$

2. To use the integral test on a series  $\sum a_n$ , the function  $f(x)$  satisfying  $f(n) = a_n$  must be positive, continuous, and decreasing. Determine whether or not the following series satisfy the conditions needed for the Integral Test:

$$\sum_{n=1}^{\infty} \frac{2n-1}{3n+1}$$

**Problems:**

1. By drawing a picture similar to the one in class or in the book that was used to justify the Integral Test, **rank the following from greatest to least**, assuming  $a_n = f(n)$ . (You may also assume that  $f(x)$  is positive, continuous, and decreasing.)

$$\int_1^6 f(x)dx, \quad \sum_{i=1}^5 a_i, \quad \sum_{i=2}^6 a_i$$

2. The error we generate when using a partial sum,  $S_n$  to estimate the sum of a series is denoted  $R_n$  for remainder.

$$R_n = S - S_n = a_{n+1} + a_{n+2} + a_{n+3} + \dots$$

Suppose that  $\sum_{n=1}^{\infty} a_n = S$ ,  $f(x)$  is continuous, decreasing, and positive on  $[1, \infty)$ , and  $f(n) = a_n$ .

- (a) Using a picture similar to the one you made in (1), justify the following formula:

$$\int_{n+1}^{\infty} f(x)dx \leq R_n \leq \int_n^{\infty} f(x)dx.$$

Conclude that

$$S_n + \int_{n+1}^{\infty} f(x)dx \leq S \leq S_n + \int_n^{\infty} f(x)dx.$$

- (b) Estimate  $\sum_{n=1}^{\infty} \frac{1}{n^2}$  to within 0.25.

3. Show that we can use the Integral Test to determine whether or not the following sums converge, and then determine whether or not they do.

(a)  $\sum_1^{\infty} \frac{\sqrt{n} + 4}{n^2} dx$

(b)  $\sum_{n=1}^{\infty} \frac{n}{n^4 + 1}$