

Math 162: Calculus IIA

Final Exam, Saturday Edition

December 12, 2020

NAME (please print legibly): _____

Your University ID Number: _____

Your University email _____

Write the name of your proctor here.

Pledge of Honesty

I affirm that I will not give or receive any unauthorized help on this exam and that all work will be my own.

Signature: _____

Instructions

- You may not consult the textbook, your notes, the internet, your classmates, friends or any other external source of information. **YOUR WEBCAM MUST BE ON AT ALL TIMES.**
- If you have access to a printer, you may print this exam and write your answers in the spaces provided. Otherwise, write the answers to each problem on a separate sheet of paper. **YOU MUST ALSO WRITE AND SIGN THE PLEDGE OF HONESTY AND GIVE ALL OF THE INFORMATION REQUESTED ABOVE.**
- Show your work and justify your answers. You may use the formulas on the next page. You may not receive full credit for a correct answer if insufficient work is shown or insufficient justification is given.
- You must finish work on this exam by 9:15, and then scan and upload it to Gradescope as previously instructed by 9:30. Exams received after that time will be subject to a penalty.

Trig formulas:

- $\cos^2(x) + \sin^2(x) = 1$
- $\sec^2(x) - \tan^2(x) = 1$
- $\sin(2x) = 2 \sin(x) \cos(x)$
- $\cos^2(x) = \frac{1 + \cos(2x)}{2}$
- $\sin^2(x) = \frac{1 - \cos(2x)}{2}$

Trigonometric substitution tricks for odd powers of secant and even powers of tangent:

- $u = \sec(\theta) + \tan(\theta)$
- $\sec(\theta)d\theta = \frac{du}{u}$
- $\sec(\theta) = \frac{u^2 + 1}{2u}$
- $\tan(\theta) = \frac{u^2 - 1}{2u}$

Integration by parts:

$$\int u dv = uv - \int v du$$

Polar coordinate formulas:

- Area:

$$\frac{1}{2} \int r^2 d\theta$$

- Arc length:

$$\int \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta$$

Parametric equation formulas:

- Newton's notation: $\dot{x} = dx/dt$ $\dot{y} = dy/dt$

- Slope of tangent line: $dy/dx = \dot{y}/\dot{x}$.

- Second derivative

$$\frac{d^2y}{dx^2} = \frac{d(\dot{y}/\dot{x})/dt}{\dot{x}}.$$

Curve is concave up/down when this is positive/negative.

- Arc length:

$$\int \sqrt{\dot{x}^2 + \dot{y}^2} dt.$$

Power series formulas:

- MACLAURIN series for $f(x)$:

$$\sum_{n=0}^{\infty} \frac{f^{(n)}(0)x^n}{n!}.$$

- Maclaurin series for specific functions:

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} \qquad \cos x = \sum_{m=0}^{\infty} \frac{(-1)^m x^{2m}}{(2m)!} \qquad \sin x = \sum_{m=0}^{\infty} \frac{(-1)^m x^{2m+1}}{(2m+1)!}$$

$$\arctan x = \sum_{m=0}^{\infty} \frac{(-1)^m x^{2m+1}}{2m+1} \qquad \ln(1+x) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1} x^n}{n}$$

$$(1+x)^k = \sum_{n=0}^{\infty} \binom{k}{n} x^n \qquad \text{where} \quad \binom{k}{n} = \frac{k(k-1)\cdots(k-n+1)}{n!}$$

- TAYLOR series for $f(x)$ about a :

$$\sum_{n=0}^{\infty} \frac{f^{(n)}(a)(x-a)^n}{n!}.$$

- The n th partial sum of the above, also called the n th Taylor polynomial, is

$$T_n(x) = \sum_{k=0}^n \frac{f^{(k)}(a)(x-a)^k}{k!},$$

and the n th Taylor remainder is $R_n(x) = f(x) - T_n(x)$. Taylor's inequality says that

$$|R_n(x)| \leq \frac{M|x-a|^{n+1}}{(n+1)!},$$

when x is in an interval centered at a in which $|f^{(n+1)}| \leq M$.

1. (25 points)

Consider the solid formed by taking the area between $y = \sqrt[3]{x}$, $y = 2$ and the y -axis, and revolving it about the y -axis.

Suppose x and y are measured in meters, and a container in the shape of the solid were filled with water of density 1000kg/m^3 from the bottom to $y = 1$ meter. How much energy (in Joules) would it take to lift the water over the top of the cup? (You can approximate gravitational acceleration by 9.8m/sec^2 .)

ANSWER:

2. (25 points)

Compute the following integral:

$$\int \frac{2x^2 - x + 4}{x^3 + 4x} dx.$$

ANSWER:


3. (25 points) Find the arc-length of the parametric curve

$$x = 2 \cos t + \cos 2t, \quad y = 2 \sin t - \sin 2t, \quad 0 \leq t \leq 2\pi.$$

by doing it for $0 \leq t \leq 2\pi/3$ and multiplying your answer by 3.

YOU MAY WANT TO USE THE TRIG IDENTITIES $\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$ AND $\sin^2 \theta = (1 - \cos 2\theta)/2$.

The curve for $0 \leq t \leq 2\pi$ is pictured below.



ANSWER:

4. (25 points)

Let $g : (0, \infty) \rightarrow \mathbb{R}$ be a positive, increasing function that satisfies

$$\lim_{x \rightarrow \infty} \frac{g(x)}{x} = 1.$$

Determine whether the series is absolutely convergent, conditionally convergent, or divergent.

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{g(n) + n}$$

ANSWER:


5. (20 points) Let r be a real number with $0 < r < 1$. Determine whether the series is absolutely convergent, conditionally convergent, or divergent.

$$\sum_{n=2}^{\infty} \frac{1}{n(\ln n)^r}.$$

ANSWER:

6. (25 points)

(10 points) (a) (10 points) Find the area inside the polar curve $r = 2\cos(\theta)$ and outside the polar curve $r = \sqrt{2}$, as shown below.



ANSWER:

(b) (10 points) Find the arc length of the boundary of the region of part (a).

ANSWER:

7. (25 points) Find the radius of convergence and interval of convergence of the series

$$\sum_{n=2}^{\infty} (-1)^n \frac{x^n}{3^n n (\ln n)^2}.$$

ANSWER:

8. (25 points)

(a) (15 points) Find the Taylor series centered at 0 of the function $\ln(1 - (x/2)^2)$, as well as the radius and interval of convergence.

ANSWER:

(b) (10 points) Write the integral

$$\int_0^x \ln(1 - (t/2)^2) dt$$

as a power series in x .

ANSWER:

