

Math 162: Calculus IIA

First Midterm Exam, Evening Edition ANSWERS

October 10, 2020

Trigonometric formulas:

- $\cos^2(x) + \sin^2(x) = 1$
- $\sin(2x) = 2 \sin(x) \cos(x)$
- $\cos^2(x) = \frac{1 + \cos(2x)}{2}$
- $\sin^2(x) = \frac{1 - \cos(2x)}{2}$

Trigonometric substitution tricks for odd powers of secant and even powers of tangent:

- $u = \sec(\theta) + \tan(\theta)$
- $\sec(\theta)d\theta = \frac{du}{u}$
- $\sec(\theta) = \frac{u^2 + 1}{2u}$
- $\tan(\theta) = \frac{u^2 - 1}{2u}$

Integration by parts:

$$\int u dv = uv - \int v du$$

1. (30 points) The problem has five parts, each worth 6 points. Each depends on having correct answers to the previous parts. You should not expect to get credit for an answer based on incorrect information.

Consider the integral

$$\int \frac{x^7 dx}{x^4 - 10x^2 + 9}.$$

- (a) Rewrite the integrand (NOT the integral, but the function being integrated) as the sum of a polynomial and a fraction in which the numerator is a polynomial of degree less than 4. SHOW YOUR WORK. You will not get credit for merely writing the correct answer.

Answer:

Here is the relevant long division of polynomials.

$$\begin{array}{r} x^3 + 10x \\ x^4 - 10x^2 + 9 \overline{) x^7} \\ \underline{-x^7 + 10x^5} \quad -9x^3 \\ 10x^5 \quad -9x^3 \\ \underline{-10x^5 + 100x^3 - 90x} \\ 91x^3 - 90x \end{array}$$

We see that the quotient is $x^3 + 10x$, and the remainder is $91x^3 - 90x$. This means that the integrand is

$$x^3 + 10x + \frac{91x^3 - 90x}{x^4 - 10x^2 + 9}.$$

- (b) Write the denominator of the fraction as a product of linear factors.

Answer:

$$x^4 - 10x^2 + 9 = (x^2 - 1)(x^2 - 9) = (x - 1)(x + 1)(x - 3)(x + 3).$$

- (c) Are there any values of x for which the integrand is not defined? If so, what are they and why?

Answer:

The denominator of the fraction is zero when x is ± 1 or ± 3 . Since division by zero is not defined, the integrand is not defined at those values of x .

- (d) Write the fraction you found in part (a) as a sum of fractions with constant numerators, which you may denote by letters of the alphabet. YOU NEED NOT FIND THE VALUES OF THESE CONSTANTS.

Answer:

$$\frac{91x^3 - 90x}{x^4 - 10x^2 + 9} = \frac{A}{x - 1} + \frac{B}{x + 1} + \frac{C}{x - 3} + \frac{D}{x + 3}.$$

- (e) Write the original integral in terms of the constants you named in part (d).

Answer:

$$\begin{aligned} \int \frac{x^7 dx}{x^4 - 10x^2 + 9} &= (x^3 + 10x) dx + \int \frac{(91x^3 - 90x) dx}{x^4 - 10x^2 + 9} \\ &= (x^3 + 10x) dx + \int \frac{A dx}{x - 1} + \int \frac{B dx}{x + 1} + \int \frac{C dx}{x - 3} + \int \frac{D dx}{x + 3} \\ &= \frac{x^4}{4} + 5x + A \ln |x - 1| + B \ln |x + 1| + C \ln |x - 3| + D \ln |x + 3| + c. \end{aligned}$$

2. (20 points) This a work problem with metric units. Assume that acceleration due to gravity is A meters per second per second. You should give your answer in joules as a multiple of $A\pi$. The density of water is a thousand kilograms per cubic meter.

Consider the region of the xy -plane bounded by the curve $y = x^2$ and the lines defined by $x = 0$ and $y = 3$. Rotate this region about the y -axis to obtain a solid region or bowl, which is filled with water. How much work is needed to pump the water about over the top of the bowl?

Answer:

We need to divide the solid region into horizontal layers, one for each value of y . Thus it is convenient to write x as a function of y , namely $x = \sqrt{y}$ for $0 \leq y \leq 3$.

The radius of such a layer is x , so its area in square meters is $\pi x^2 = \pi y$, and its volume is therefore $\pi y dy$ cubic meters. This means its mass is $1000\pi y dy$ kilograms, so the gravitational force acting on it is $1000A\pi y dy$ newtons. The distance to the top of the solid is $(3 - y)$ meters, so the work needed to lift it is $1000A\pi y(3 - y) dy$ joules.

Thus the total amount of work in joules is

$$\int_0^3 1000A\pi y(3 - y) dy = 1000A\pi \int_0^3 (3y - y^2) dy$$

$$\begin{aligned}
 &= 1000A\pi \left(\frac{3y^2}{2} - \frac{y^3}{3} \right) \Big|_0^3 \\
 &= 1000A\pi \left(\frac{27}{2} - \frac{27}{3} \right) \\
 &= 4500A\pi.
 \end{aligned}$$

3. (30 points)

(a) (15 points) Use integration by parts twice to find a formula for

$$\int x^n e^{-x^2} dx \quad \text{in terms of} \quad \int x^{n-4} e^{-x^2} dx \quad \text{for } n > 4.$$

Hint: start with $x^n e^{-x^2} = \left(-\frac{1}{2}x^{n-1}\right) \left(-2xe^{-x^2}\right)$, then use

$$u = -\frac{1}{2}x^{n-1}, \quad dv = -2xe^{-x^2} dx.$$

Answer:

Using integration by parts twice we have

$$\begin{aligned}
 \int x^n e^{-x^2} dx &= \int \left(-\frac{1}{2}x^{n-1}\right) \left(-2xe^{-x^2}\right) dx \\
 &= -\frac{1}{2} \int x^{n-1} \left(e^{-x^2}\right)' dx \\
 &= -\frac{1}{2} \left(x^{n-1} e^{-x^2} - \int (n-1)x^{n-2} e^{-x^2} dx \right) \\
 &= -\frac{1}{2} x^{n-1} e^{-x^2} + \frac{n-1}{2} \int x^{n-2} e^{-x^2} dx \\
 &= -\frac{1}{2} x^{n-1} e^{-x^2} + \frac{n-1}{2} \int \left(-\frac{1}{2}x^{n-3}\right) \left(e^{-x^2}\right)' dx \\
 &= -\frac{1}{2} x^{n-1} e^{-x^2} - \frac{n-1}{4} \left(x^{n-3} e^{-x^2} - \int (n-3)x^{n-4} e^{-x^2} dx \right) \\
 &= -\frac{1}{2} x^{n-1} e^{-x^2} - \frac{n-1}{4} x^{n-3} e^{-x^2} + \frac{(n-1)(n-3)}{4} \int x^{n-4} e^{-x^2} dx
 \end{aligned}$$

(b) (15 points) Let

$$I_n = \int x^n e^{-x^2} dx.$$

Use your formula of part (a) to find I_9 . You should not expect to get credit for an answer based on an incorrect formula.

Answer:

$$\begin{aligned} I_9 &= \int x^9 e^{-x^2} dx \\ &= -\frac{1}{2}x^8 e^{-x^2} - \frac{8}{4}x^6 e^{-x^2} + \frac{8 \cdot 6}{4} \int x^5 e^{-x^2} dx \\ &= -\frac{1}{2}x^8 e^{-x^2} - 2x^6 e^{-x^2} + 12 \left(-\frac{1}{2}x^4 e^{-x^2} - \frac{4}{4}x^2 e^{-x^2} + \frac{4 \cdot 2}{4} \int x e^{-x^2} dx \right) \\ &= -\frac{1}{2}x^8 e^{-x^2} - 2x^6 e^{-x^2} - 6x^4 e^{-x^2} - 12x^2 e^{-x^2} + 24 \int x e^{-x^2} dx \\ &= -\frac{1}{2}x^8 e^{-x^2} - 2x^6 e^{-x^2} - 6x^4 e^{-x^2} - 12x^2 e^{-x^2} - 12e^{-x^2} + C \end{aligned}$$

4. (20 points)

Compute the following integral:

$$\int \frac{\sqrt{4-x^2}}{x^4} dx$$

Answer:

Let $\cos(\theta) = x/2$. So $dx = -\sin(\theta)/2d\theta$. $\sqrt{4-x^2} = \sin(\theta)$.

After trig substitution, the integral is

$$-\int \frac{\sin^2(\theta)}{4 \cos^4(\theta)} d\theta.$$

Rewritten, this is

$$-\frac{1}{4} \int \tan^2(\theta) \sec^2(\theta) d\theta.$$

Set $u = \tan(\theta)$, so $du = \sec^2(\theta)d\theta$. After this substitution the integral is

$$-\frac{1}{4} \int u^2 du.$$

This equals

$$-\frac{1}{12}u^3 + C.$$

Substituting back $u = \tan(\theta)$ we get

$$-\frac{1}{12}\tan^3(\theta) + C.$$

In terms of x , $\tan(\theta) = \frac{\sqrt{4-x^2}}{x}$, so we get

$$-\frac{(4-x^2)^{3/2}}{12x^3} + C.$$