

Math 162 second midterm exam
November 11, 2021

HANDY DANDY FORMULAS

Integration by parts formula:

$$\int u dv = uv - \int v du$$

Trigonometric identities:

$$\begin{aligned} \cos^2 \theta + \sin^2 \theta &= 1 & \sec^2 \theta - \tan^2 \theta &= 1 \\ \sin(\alpha + \beta) &= \sin \alpha \cos \beta + \cos \alpha \sin \beta & \cos(\alpha + \beta) &= \cos \alpha \cos \beta - \sin \alpha \sin \beta \\ \cos^2 \theta &= \frac{1 + \cos 2\theta}{2} & \sin^2 \theta &= \frac{1 - \cos 2\theta}{2} \end{aligned}$$

Derivatives of trig functions.

$$\begin{aligned} \frac{d \sin x}{dx} &= \cos x & \frac{d \tan x}{dx} &= \sec^2 x & \frac{d \sec x}{dx} &= \sec x \tan x \\ \frac{d \cos x}{dx} &= -\sin x & \frac{d \cot x}{dx} &= -\csc^2 x & \frac{d \csc x}{dx} &= -\csc x \cot x \end{aligned}$$

Trigonometric substitution for integrals of the form

$$\int \tan^m x \sec^n x dx \quad \text{with } n > 0,$$

known in Doug's section as *the rabbit trick*.

$$\begin{aligned} u &= \sec x + \tan x & \sec x dx &= \frac{du}{u} \\ \sec x &= \frac{u^2 + 1}{2u} & \tan x &= \frac{u^2 - 1}{2u} \end{aligned}$$

Area of surface of revolution in rectangular coordinates, $y = f(x)$ with $a \leq x \leq b$

- about the x -axis: $S = 2\pi \int_a^b f(x) \sqrt{1 + f'(x)^2} dx$
- about the y -axis: $S = 2\pi \int_a^b x \sqrt{1 + f'(x)^2} dx$

MORE FORMULAS FOR YOUR ENJOYMENT

Polar coordinates

$$\begin{aligned} r &= \sqrt{x^2 + y^2} & \theta &= \arctan(y/x) & \text{for } x > 0 \\ \pi + \arctan(y/x) & \text{for } x < 0 \\ \pi/2 & \text{for } x = 0 \text{ and } y > 0 \\ 3\pi/2 & \text{for } x = 0 \text{ and } y < 0 \\ \text{undefined} & \text{for } (x, y) = (0, 0) \\ x &= r \cos \theta & y &= r \sin \theta \end{aligned}$$

Changing θ by any multiple of 2π does not change the location of the point. Changing the sign of r is equivalent to adding π to θ , which is the same as moving the point to one in the opposite direction and the same distance from the origin.

Area in polar coordinates for $r = f(\theta)$ with $\alpha \leq \theta \leq \beta$:

$$A = \int_{\alpha}^{\beta} \frac{r^2}{2} d\theta$$

Arc length formulas

- Rectangular coordinates, $y = f(x)$ with $a \leq x \leq b$:

$$S = \int_a^b \sqrt{1 + f'(x)^2} dx$$

- Polar coordinates, $r = f(\theta)$ with $\alpha \leq \theta \leq \beta$:

$$S = \int_{\alpha}^{\beta} \sqrt{r^2 + f'(\theta)^2} d\theta$$

- Parametric equations, $x = x(t)$ and $y = y(t)$ with $a \leq t \leq b$:

$$S = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$