

Math 164: Multidimensional Calculus

Midterm 2

November 17, 2015

NAME (please print legibly): _____

Your University ID Number: _____

Indicate your instructor with a check in the appropriate box:

Bobkova	TR 12:30-1:45pm	<input type="checkbox"/>
Chen	MW 3:25-4:40pm	<input type="checkbox"/>
Dummit	TR 3:25-4:40pm	<input type="checkbox"/>
Salur	MWF 09:00-09:50am	<input type="checkbox"/>

- You are responsible for checking that this exam has all 9 pages.
- No calculators, phones, electronic devices, books, notes are allowed during the exam.
- Show all work and justify all answers.
- Please sign the pledge below.

Pledge of Honesty

I affirm that I will not give or receive any unauthorized help on this exam, and that all work will be my own.

Signature: _____

QUESTION	VALUE	SCORE
1	8	
2	8	
3	12	
4	15	
5	14	
6	14	
7	14	
8	15	
TOTAL	100	

1. (8 points) Suppose the equation $x^2z^4 + 2ye^{x+z} = 5$ defines z implicitly as a function of x and y . Find the value of $\frac{\partial z}{\partial x}$ at the point $(x, y, z) = (-1, 2, 1)$.

2. (8 points) Find an equation for the tangent plane to the surface $xy^2z + \ln(x+2y+z) = 2$ at the point $(x, y, z) = (2, -1, 1)$.

3. (12 points) Consider the function $f(x, y) = xy^2$ and the point $P(1, 2)$.

(a) Find the rate of change of f at P in the direction of the vector $\mathbf{u} = \left\langle \frac{2}{\sqrt{13}}, \frac{3}{\sqrt{13}} \right\rangle$.

(b) Find the maximal rate of change of f at P and the direction in which it occurs.

4. (15 points) Find the absolute minimum and maximum values of the function $f(x, y) = 4x + 6y - x^2 - y^2$ on the region $D = \{(x, y) : 0 \leq x \leq 4, 0 \leq y \leq 5\}$.

5. (14 points) Find the greatest and smallest values of $f(x, y) = 2xy$ on the ellipse

$$g(x, y) = \frac{x^2}{4} + y^2 = 1 = k. \quad g(x, y) = k \text{ is closed \& bounded, } f \text{ cont's } \Rightarrow f$$

LM: attains ~~or~~ max and min on $g(x, y) = k$. These extrema of f along the level curve $g(x, y) = k$ will occur when the tangent line to $g(x, y) = k$ is parallel to the tangent line to a level curve of f , or when ∇f is \parallel to ∇g , i.e. $\nabla f = \lambda \nabla g$.

$$\nabla f = \langle 2y, 2x \rangle \text{ and } \nabla g = \langle \frac{x}{2}, 2y \rangle$$

$$\text{so } \nabla f = \lambda \nabla g \Rightarrow \begin{cases} 2y = \lambda \frac{x}{2} \\ 2x = \lambda 2y \end{cases} \text{ so we must solve } \begin{cases} \textcircled{1} 4y - \lambda x = 0 \\ \textcircled{2} 2x - 2\lambda y = 0 \\ \textcircled{3} x^2 + 4y^2 = 4 \end{cases}$$

Algebra: By $\textcircled{1}$, either $x=0$ or $\lambda = \frac{4y}{x}$. If $\underline{x=0}$, by $\textcircled{3}$ we get $y = \pm 1$, so $\boxed{(0, \pm 1)}$

$$\text{If } \underline{\lambda = \frac{4y}{x}}, \text{ by } \textcircled{2} \quad 2x - 2\left(\frac{4y}{x}\right)y = 0, \text{ or } 2x^2 - 8y^2 = 0.$$

$$\text{By } \textcircled{3}, \quad 2x^2 + 8y^2 = 8, \text{ so we have } \begin{aligned} 2x^2 - 8y^2 &= 0 \\ \text{and } 2x^2 + 8y^2 &= 8 \end{aligned}$$

$$\text{Adding, we get } 4x^2 = 8. \text{ Subtracting, we get } \begin{aligned} 16y^2 &= 8 \\ y^2 &= \frac{1}{2} \end{aligned}$$

$$\begin{aligned} x^2 &= 2 \\ \boxed{x = \pm\sqrt{2}} \end{aligned}$$

$$\boxed{y = \pm\sqrt{\frac{1}{2}}}$$

Check values:

$$f(0, \pm 1) = 0$$

$$f(\pm\sqrt{2}, \pm\sqrt{\frac{1}{2}}) = 2 \leftarrow \text{abs. max of } f \text{ on } g=k$$

$$f(\pm\sqrt{2}, \mp\sqrt{\frac{1}{2}}) = -2 \leftarrow \text{abs. min. of } f \text{ on } g=k$$

6. (14 points) Evaluate the double integral

$$I = \int_0^1 \int_0^2 xye^{xy^2} dx dy.$$


$$I = \int_0^2 \int_0^1 xye^{xy^2} dy dx \quad \text{by Fubini since } f(x,y) = xye^{xy^2} \text{ con'ts}$$

$$= \int_0^2 \left[\frac{1}{2} e^{xy^2} \right]_{y=0}^1 dx = \int_0^2 \left(\frac{1}{2} e^0 - \frac{1}{2} e^0 \right) dx = \frac{1}{2} x - \frac{1}{2} e^x \Big|_0^2$$

\uparrow
 $u = xy^2$
 $\frac{1}{2} du = xy dy$

$$= 1 - \frac{1}{2} e^2 - \left(0 - \frac{1}{2} \right) = \boxed{\frac{3}{2} - \frac{1}{2} e^2}$$

7. (14 points) Evaluate $\iint_R y \, dA$ where R is the triangular region with vertices $(0,0)$, $(3,0)$, and $(1,1)$.




$$R = \{(x,y) \mid 0 \leq y \leq 1, y \leq x \leq 2y-3\}$$

$$I = \int_0^1 \int_y^{2y-3} y \, dx \, dy = \int_0^1 [xy]_y^{2y-3} \, dy$$

$$= \int_0^1 ((2y-3)y - y^2) \, dy = \int_0^1 (y^2 - 3y) \, dy = \frac{1}{3}y^3 - \frac{3}{2}y^2 \Big|_0^1 = \frac{1}{3} - \frac{3}{2} \boxed{\frac{7}{6}}$$

$$\boxed{I = -\frac{7}{6}}$$

8. (15 points) Consider the integral $I = \iint_R (x^2 + y^2)^4 dA$ where R is the region above the line $y = 0$, below the line $y = x$, and inside the circle $x^2 + y^2 = 4$, as pictured below:



- (a) Set up (no need to evaluate) an iterated double integral for I in rectangular xy -coordinates with your choice of integration order.

$$R = \{(x, y) \mid 0 \leq y \leq \sqrt{2}, y \leq x \leq \sqrt{4-y^2}\}$$

$$I = \int_0^{\sqrt{2}} \int_y^{\sqrt{4-y^2}} (x^2 + y^2)^4 dx dy$$

- (b) Evaluate the double integral in polar $r\theta$ -coordinates.

$$R = \{(r, \theta) \mid 0 \leq r \leq 2, 0 \leq \theta \leq \frac{\pi}{4}\}$$

$$f(r \cos \theta, r \sin \theta) = (r^2)^4 = r^8$$

$$dA = r dr d\theta$$

$$I = \int_0^{\pi/4} \int_0^2 r^8 r dr d\theta = \int_0^{\pi/4} \frac{1}{10} r^{10} \Big|_0^2 d\theta = \frac{2^{10}}{10} \pi/4 = \boxed{\frac{2^{10} \pi}{40}}$$